

Depitox

A soluble concentrate containing 500 g/l (42% w/w) 2,4-D as the dimethylamine salt.

For the selective control of the annual and perennial broad-leaved weeds in winter and spring wheat, barley and rye, winter oats, established agricultural and amenity grassland, managed amenity turf, apple and pear orchard floors and for control of aquatic weeds.

PRODUCT BENEFITS

- Controls many important broad-leaved weeds in cereals including Charlock, Common pansy, Fat hen and Shepherd's purse.
- Use on grass floors under orchards.
- Can be used in a wide range of cereal crops.
- Controls many key weeds in grassland.
- Tank mix recommendation for Ragwort control.

LERAP category: Unclassified

Pack size: 10 litres

Storage: PROTECT FROM FROST



IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL/AQUATIC HERBICIDE.

| Crop | Maximum individual dose | Maximum total dose | Latest time of application |
|--|--------------------------------|---------------------------|------------------------------------|
| Winter wheat, winter and spring rye | 2.5 l/ha | 2.5 l/ha per crop | before first node detectable stage |
| Spring wheat, winter and spring barley, winter oats | 2.0 l/ha | 2.0 l/ha per crop | |
| Winter and spring wheat, winter and spring barley, winter oats, winter and spring rye (undersown with grass) | 1.0 l/ha | 1.0 l/ha per crop | |
| Grassland | 3.3 l/ha | 3.3 l/ha per year | before the crop is 25cm high |
| Apple (around), Pear (around) | 2.8 l/ha | 2.8 l/ha per year | – |
| Amenity grassland, managed amenity turf | 3.3 l/ha | 9.9 l/ha per year | |
| Enclosed water, open waters, Land immediately adjacent to aquatic areas | 9.0 l/ha | 9.0 l/ha per year | |

Other specific restrictions:

- When applying in aquatic situations a maximum concentration of 9.0 litres product per 250 litres spray solution must not be exceeded.
- Livestock must be kept out of treated areas for at least two weeks following treatment and until poisonous weeds such as ragwort have died and become unpalatable.
- Users must consult the appropriate water regulatory body (Environment Agency/Scottish Environment Protection Agency) before using the product near water and must obtain their agreement before using this product to control aquatic weeds.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

MAPP 13258

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS

DO NOT use DEPITOX on the seedbed before sowing any crop.

DO NOT sow any crop into soil treated with DEPITOX for at least three months after application.

DO NOT graze grass for at least 14 days after spraying.

DO NOT mow or roll four days before or after application. The first four mowings after treatment must be composted for at least six months before use.

DO NOT treat newly established grass or turf less than one year old.

DO NOT treat grass or turf suffering from stress caused by drought, frost, disease or other adverse factors.

DO NOT roll or harrow crops for seven days either before or after application of DEPITOX.

Cautions to use in or near water courses

Under the Water Resources Act (1991) and the Control of Pollution Act (1974), the Environmental Agency in England and Wales and the Scottish Environment Protection Agency in Scotland are legally charged with the prevention of pollution in rivers. If the water which it proposed to treat with DEPITOX is subject to the above legislation (e.g. enters a river or is part of a river system), consult the Environment Agency or Scottish Environment Protection Agency before treatment. For Northern Ireland the Environment and Heritage Services holds similar responsibilities to the Environment Agency and Scottish Protection Agency.

The official '*Guidelines for the Use of Herbicides on Weeds in or near Watercourses and Lakes*' must be read before use of DEPITOX in the listed aquatic situations. Copies are available from DEFRA Publications, Admail 6000, London SW1A 2XX (telephone 08459 556000) quoting Ref: PB 2289.

Water containing the herbicide must not be used for irrigation purposes within three weeks of treatment, or until the concentration in water is below 0.5 ppm.

WEEDS CONTROLLED

Apply when the majority of annual weeds are at the seedling* stage. For the control of perennial weeds in established grassland, the best results are obtained if spraying is carried out shortly before flowering. Whilst spraying at this late stage will not give complete control of annual weeds, it may effectively check most of the species mentioned. A second application may be necessary to provide an adequate level of weed control on amenity grassland and managed amenity turf.

* Seedling = Fully expanded cotyledons to two expanded true leaves.

WEEDS CONTROLLED – Cereals

| Weed | Dose | Notes |
|--|-------------|--|
| Charlock Mustard, black | 700 ml | These weeds will be completely or almost completely killed when applications are made in the cotyledon to early flower-bud stage |
| Fat hen Mustard, treacle Mustard, white Pennycress, field Tare, hairy | 1.4 l/ha | These weeds will be completely or almost completely killed when applications are made in the cotyledon to early flower-bud stage |
| Buttercup, corn Nettle, small Radish, wild (Runch) Shepherd's purse | | These weeds will be completely or almost completely killed when applications are made in the cotyledon up to two leaf stage or moderately susceptible at four leaves to early flower-bud stage |
| Forget-me-not, field Orache, common Poppy, common Sowthistle, prickly Sowthistle, smooth Turnip, wild | | These weeds will be moderately susceptible (with or without mortality) when applications are made in the cotyledon up to two leaf stage |
| Bindweed, black Bugloss Bugloss, viper's Chickweed, common Crane's-bill, dove's-foot Fumitory, common Gromwell, field Groundsel Knotgrass Mouse-ear, common Nightshade, black Persicaria, pale Pimpernel, scarlet Redshank Shepherd's needle Speedwell, common field Speedwell, green field Speedwell, ivy-leaved Speedwell, wall Spurge, sun | | These weeds will be moderately resistant when applications are made in the cotyledon up to two leaf stage |

Cereals *continued*

| Weed | Dose | Notes |
|---|--------------|---|
| Orache, common Poppy, common Sowthistle, smooth | 2.0 l/ha | These weeds will be susceptible when applications are made in the cotyledon up to four leaf stage or moderately resistant at six leaves to early flower-bud stage |
| Knotgrass Mayweed, scentless | 2.0 l/ha | These weeds will be moderately resistant when applications are made in the cotyledon up to two leaf stage or resistant at four leaves to early flower-bud stage |
| Thistle, creeping* | 2.0–2.5 l/ha | These weeds will be susceptible when applications are made in the cotyledon up to early flower-bud stage |

* Aerial growth only.

Hoary cress – Good control of this perennial weed can be achieved by treatment in winter cereal crops over two successive seasons using 1.6–1.8 l/ha dose of DEPITOX. Apply after the shoots are 25–150 mm high, up to but before flowering.

Amenity grassland and managed amenity turf

| Weed | Dose | Notes |
|--|----------|---|
| Buttercup, creeping Hawkweed, mouse-ear Plantains Thrift | 2.8 l/ha | These weeds are consistently killed by one application |
| Bedstraw, heath Buttercup, bulbous Catsear Chickweed, common Daisy Dandelion Dock, curled Hawkbit, rough Hawk's-beard, smooth Pennywort, marsh Sea, milkwort Sorrel, common Sorrel, sheep's Stork's-bill, common Stork's-bill, sea Thistle, dwarf | | Sometimes killed by one application but may require a second treatment to give complete control |

Continued...

WEEDS CONTROLLED – Amenity grassland and managed amenity turf *continued*

| Weed | Dose | Notes |
|--|----------|--|
| Celandine, lesser Mouse-ear, common Pearlwort, procumbent Self heal Yarrow | | Some effect from one application, but two applications required to give a useful level of control |
| Ragwort, common* | 3.3 l/ha | Moderately susceptible. Sometimes killed by one application but may require further treatment to give complete control |

* treatment will normally kill plants at all stages of growth up to early bud stage. For best levels of control, treat in April–June when rosettes are growing strongly but before flower buds are well formed.

Agricultural grassland (including grass floors under apple and pear trees)

| Weed | Dose | Notes |
|--|----------|--|
| Buttercup, creeping Hawkbit, autumn Hawk's-beard, rough Plantain, greater Plantain, hoary Plantain, ribwort Sandwort, thyme-leaved | 2.8 l/ha | These weeds are susceptible at all stages of growth up to the beginning of flowering with good control of shoots and roots in established plants |
| Buttercup, bulbous Dock, broad-leaved | | Seedlings and shoots are susceptible but established plants in grassland will not be controlled |
| Dandelion Dock, curled Nettle, common Rush, soft * Thistle, creeping | | Seedlings and shoots are susceptible but only aerial growth of established plants is usually controlled |
| Thistle, spear | | Seedlings are susceptible but only aerial growth of established plants is usually controlled |
| Bartsia, red Bindweed, hedge Burdock, lesser Buttercup, meadow Catsear Chicory Cress, hoary Daisy | | These weeds are well controlled in the seedling or shoot stage with useful suppression or death of aerial parts at later growth stages |

Agricultural grassland (including grass floors under apple and pear trees) *continued*

| Weed | Dose | Notes |
|-----------------------------|-----------------------|--|
| Dock, clustered | 2.8 l/ha | These weeds are well controlled in the seedling or shoot stage with useful suppression or death of aerial parts at later growth stages |
| Fleabane, common | | |
| Goatsbeard | | |
| Hawkbit, rough | | |
| Hawksbeard, smooth | | |
| Hawkweed, mouse-ear | | |
| Hemp-nettle, large flowered | | |
| Knapweed, common | | |
| Knawel, annual | | |
| Loosestrife, purple | | |
| Mugwort | | |
| Ox-tongue, bristly | | |
| Plantain, buck's horn | | |
| Radish, horse | | |
| Scabious, field | | |
| Self-heal | | |
| Thistle, musk | | |
| Thorn apple | | |
| Vetch, common | | |
| Vetch, tufted | | |
| Horsetail, field ** | 2.8 l/ha | Only controls shoots which are well developed (preferably about 30 cm high). Control of established plants is variable. Re-growth will occur in following season |
| Horsetail, marsh ** | | |
| Sorrel, common | | |
| Sorrel, sheep's | 2.0 l/ha | Provides useful control of shoots only |
| Sowthistle, perennial | | |
| Bindweed, field**** | 3.3 l/ha ⁺ | Moderately susceptible. Aerial growth usually killed and a useful measure of long term control obtained under suitable conditions |
| Ragwort, common*** | | |

* May be controlled by application in April to June when growing well. For best results, cut the rushes four weeks after treatment or cut them four weeks before application and remove stems before spraying.

** Use 2.8 l/ha and spray when growing well in May or early June. Top growth is removed or considerably reduced for the season of treatment. In grassland for hay or silage, shoot kill may be obtained by using 2.0 l/ha two weeks before cutting.

*** Treatment will normally kill plants at all stages of growth up to early bud stage. For best levels of control, treat in April–June when rosettes are growing strongly but before flower buds are well formed.

**** In order to obtain maximum effect in the year after treatment, spraying should be delayed until shoots are well developed.

+ Application rate of 3.3 l/ha is not permissible around apple and pear trees for the control of Common ragwort, although a maximum individual dose and maximum total dose of 2.8 l/ha is permitted.

Aquatic uses

| Weed | Dose | Notes |
|---|----------|--|
| Mint, water Plantain, water | 4.4 l/ha | Complete or almost complete kill in year of application |
| Cress, water Rush, soft | 9.0 l/ha | Complete or almost complete kill in year of application |
| Bur-reed, branched Willowherb, great | | Partial kill and effective suppression in year of treatment |
| Bistort, amphibious Horsetail, water | | Temporary suppression. Re-growth occurring during year of treatment. |

Ragwort control

Ragwort is an injurious weed and those who permit it to grow unchecked on their land are liable to prosecution under the Weed Act (1959).

Dose

Agricultural grassland: DEPITOX at 2.8 l/ha + Agritox 50 (MAFF 07400) at 2.0 l/ha.

Do not apply 2.8 l/ha DEPITOX alone as this will not give reliable control of Ragwort.

Timing

Agricultural grassland (including grass floors under apple and pear trees), amenity grassland and managed amenity turf.

Spray when the majority of plants are in the rosette stage and growing vigorously in the autumn or spring but before the flower spines start to grow. DEPITOX should be applied in good growing conditions. Treatment of Ragwort should always be part of a programme and repeat application may be necessary together with removal of any flower heads in the summer to reduce seed return to the soil. Fields for hay or silage the following spring should be sprayed in the preceding autumn. Fields to be grazed should be treated in the spring.

NOTE: It is important that all livestock are kept out of treated areas for at least two weeks following treatment and until the Ragwort has died and become unpalatable.

CROP SPECIFIC INFORMATION

| Crop | Dose | Maximum total Dose | Timing and notes |
|--|----------------|--------------------|---|
| Winter Wheat, Winter and Spring Rye | 0.7–2.5 l/ha | 2.5 per crop | Winter cereals: Apply in the spring from the leaf sheath erect stage but before the first node detectable stage Spring cereals: Apply from the five leaf fully expanded stage but before the first node detectable stage |
| Winter and spring barley, winter oats, spring wheat | 0.7 – 2.0 l/ha | 2.0 l/ha per crop | Winter cereals: Apply in the spring from the leaf sheath erect stage but before the first node detectable stage Spring cereals: Apply from the five leaf fully expanded stage but before the first node detectable stage |
| Winter and spring wheat, winter and spring barley, winter oats and winter and spring rye (undersown with grass) | 1.0 l/ha | 1.0 l/ha per crop | Apply in the spring following the same recommendations as for cereals. DO NOT spray with DEPITOX before under sowing. Experience has shown that when weeds and cereals form a canopy undersown crops may be safely treated using not more than 1.0 l/ha at low volume. |
| Rotational and permanent grassland established for at least one year. Do not use where clovers are an important part of the sward | 3.3 l/ha | 3.3 l/ha per year | Apply in spring to autumn at the optimum timing when grass density is low, such as after cutting or grazing, but when weeds are at a susceptible stage. Grassland may be treated with 2.8–3.3 l/ha of DEPITOX according to the weeds present. Recommended rates are given in the weed susceptibility table for grassland |

Continued

CROP SPECIFIC INFORMATION *continued*

| Crop | Dose | Maximum total Dose | Timing and notes |
|---|--------------|--------------------|---|
| Grass floors under apple and pear orchards. The orchards must have established for at least one year. Do not apply directly to trees | 2.8 l/ha | 2.8 l/ha per year | Apply in spring or autumn when weeds are actively growing. Do not spray during blossom or whilst weeds are in flower. Use low pressure nozzles to avoid spray drift. Bramley Seedling, Emneth Early and Miller's Seedling are particularly susceptible to spray drift. Pears are more susceptible to spray drift than apples and are particularly susceptible to damage via root uptake |
| Amenity grassland and managed amenity turf (established for at least one year). | 3.3 l/ha | 9.9 l/ha per year | Apply in spring/summer or autumn when the growing conditions are favourable. Amenity grassland and managed amenity turf may be treated with 2.8–3.3 l/ha of DPEITOX. The expected levels of control are detailed in the weed susceptibility table for amenity uses. Some perennial weeds will need subsequent application in order to achieve adequate control. A follow up application may also be needed where new seedling weeds appear. An interval of four–six weeks should elapse between applications. Clovers will receive a check. Top dressing ten days before treatment is recommended to assist kill of weeds and subsequent recovery of the sward |
| Aquatic use | 4.4–9.0 l/ha | 9.0 l/ha per year | Apply to foliage and aerial parts of aquatic weeds from May to September |

See under 'Weeds Controlled' for specific application rates for individual weeds.

DEPITOX may be used on all varieties of the listed crops within the recommended growth stages. DO NOT treat barley intended for malting or any cereal mixture with peas or beans or other legumes.

Apply in at least 110 litres of water per hectare. In grassland and turf, where weeds might be shielded by grasses, use 400 l/ha water. Refer to the table for special situation pertaining to grass floors under apples and pears. For aquatic uses apply in 250–700 litres of water per hectare.

MIXING AND SPRAYING

Before use ensure that the spraying equipment has been thoroughly cleaned. Half-fill the spray tank with clean water. With the contents of spray tank under re-circulation, add the measured quantities of DEPITOX through the filter. Top up the tank with water to the required level and maintain re-circulation until the tank is sprayed out.

Apply the recommended quantity of DEPITOX through a conventional hydraulic sprayer using a MEDIUM spray to cover the weed leaves evenly and thoroughly.

Avoid spray drift onto neighbouring crops and all broad-leaved plants outside the target area. Do not spray in windy weather. Beets, all brassicas (including oilseed rape, Swedes and turnip) lettuce, sunflowers, onions, peas, potatoes, tomatoes, cucumbers, all fruit crops (including vines) and ornamentals are particularly susceptible to 2,4-D and may be damaged by spray drift.

After each days use, wash out with water and wetting agent. Wash out again with water, drain and allow to dry. Traces of herbicide left in the sprayer may damage susceptible crops if these are subsequently sprayed using the same equipment.

WEATHER AND GROWING CONDITIONS

Apply to a dry crop when rain is not forecast for at least 12 hours. Optimum results are obtained when the weeds are actively growing under good soil and weather conditions. Reduced weed control may be obtained during drought or cold weather. If rain falls shortly after application, the effect of DEPITOX may be reduced.

RESISTANCE MANAGEMENT

When herbicides with the same mode of action are used repeatedly over several years in the same field, selection of resistant biotypes can take place. These can propagate and may become dominating. A weed species is considered to be resistant to a herbicide if it survives a correctly applied treatment at the recommended dose. A strategy for preventing and managing such resistance should be adopted. This should include integrating herbicides with a programme of cultural control measures. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

DEPITOX

A soluble concentrate containing 500 g/l (42% w/w) 2,4-D as the dimethylamine salt.

**IRRITANT**

IRRITATING TO SKIN.

RISK OF SERIOUS DAMAGE TO EYES.

HARMFUL TO AQUATIC ORGANISMS, MAY CAUSE LONG TERM ADVERSE EFFECTS IN THE AQUATIC ENVIRONMENT.

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY WITH PLENTY OF WATER AND SEEK MEDICAL ADVICE.

WEAR SUITABLE GLOVES.

WEAR SUITABLE EYE/FACE PROTECTION.

THIS MATERIAL AND ITS CONTAINER MUST BE DISPOSED OF IN A SAFE WAY.

USE APPROPRIATE CONTAINMENT TO AVOID ENVIRONMENTAL CONTAMINATION.

Contains 2,4-D. May produce an allergic reaction.

To avoid risks to man and the environment, comply with the instructions for use.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when applying by hand held equipment.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before meals and after work.

WASH CONCENTRATE from skin or eyes immediately.

Environmental protection

Do not contaminate water with the product or its container except as directed for aquatic use (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

The maximum concentration of active ingredient in treated water must not exceed 1 ppm or such lower concentration as the appropriate water regulatory authority may require.

KEEP LIVESTOCK out of treated areas for at least two weeks and until poisonous weeds such as ragwort have died and become unpalatable.

Users must consult the appropriate water/regulatory body (Environment Agency/Scottish Environment Protection Agency) before using the product near water and must obtain their agreement before using this product to control aquatic weeds.

Storage and disposal

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDSTUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, and dispose of safely.