

Legolas

Contains 300 g/l 2,4-D, 120 g/l dicamba and 400 g/l dichlorprop-p as the dimethylamine salts formulated as a soluble concentrate.

A selective weedkiller for the control of docks in grassland.

PRODUCT BENEFITS

- Control of the two major grassland weeds Docks and Creeping thistle.
- One of few products which will control Dandelion in amenity grassland.
- Unique combination of actives for grassland.

LERAP category: Unclassified

Pack size: 10 litres

Storage: PROTECT FROM FROST



IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE.

Crop	Maximum individual dose	Maximum number of treatments	Latest time of application
Agricultural grassland	2.0 l/ha	One per year	–
Amenity grassland	2.0 l/ha	Two per year	–

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

MAPP 14439

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS

LEGOLAS will cause severe damage to clover. DO NOT USE LEGOLAS on grass swards where clover is an important part of the vegetation.

DO NOT mow or roll four days before or after application. The first four mowings after treatment must be composted for at least six months before use.

DO NOT treat newly established grass less than one year old.

DO NOT treat grass crops intended for seed production.

DO NOT treat grass suffering from drought, disease or other adverse factors.

DO NOT roll or harrow crops for seven days either before or after application of LEGOLAS.

DO NOT apply when grassland is flowering.

WEEDS CONTROLLED

Weeds are most susceptible to LEGOLAS when growing vigorously under warm moist conditions. Results from spraying under cold or frosty conditions will not be as satisfactory.

Weeds	Susceptibility	Maximum growth stage
Dandelion	S	rosette (before flowering)
Dock species	S	25 cm tall or 25 cm diameter
Sowthistle, perennial	MS	rosette (before flowering)
Thistle, creeping	S ¹	rosette (before flowering)

S = Susceptible up to the growth stage listed under good growing conditions.

S¹ = A second application may be required to treat re-growth in severe infestations.

MS = Moderately susceptible.

CROP SPECIFIC INFORMATION

Established agricultural grassland

Maximum individual dose	Maximum number of applications	Water volume
2.0 l/ha	One per year	200 litres of water per hectare

Amenity grassland

Maximum individual dose	Maximum number of applications	Water volume
2.0 l/ha	Two per year	200 litres of water per hectare

Spray in the spring when weed growth is vigorous. This is usually in April, but can be carried out earlier if conditions are suitable. Best results are from two applications made in the spring of succeeding years.

Where grassland has been grazed or cut, allow at least 14 days to elapse and ensure that the weeds have re grown before treatment. Weed growth and control is usually enhanced by an application of nitrogen fertilizer two-three weeks before treatment with LEGOLAS.

SUCCEEDING CROPS

Where re-seeding is necessary do not do it within three months following treatment.

ADJACENT CROPS

DO NOT spray in windy or very still conditions. Spray drift must be avoided especially near susceptible crops such as tomatoes, lettuce, oilseed rape, vegetables, turnips, swedes, sugar beet, peas, glasshouse crops, fruit and ornamentals, etc. Avoid drift onto all broad-leaved plants outside the target area

MIXING AND SPRAYING

Half fill the spray tank with clean water and start the agitation. Pour in the required amount of LEGOLAS. Add the remainder of the water and continue agitation until spraying is completed.

WASH EQUIPMENT thoroughly immediately after use. Fill the tank with clean water and leave overnight. Spray out before storage or using other products. Traces of the product may cause damage to susceptible crops sprayed later.

COMPATIBILITY

When tank mixes are to be used, each product should be added separately to the spray tank, taking due note of any instructions given as to the order of mixing. Whilst most pesticides are compatible with LEGOLAS, check the other manufacturer's recommendations before tank mixing.

LEGOLAS

A soluble concentrate containing 300 g/l 2,4-D, 120 g/l dicamba and 400 g/l dichlorprop-p.



HARMFUL



DANGEROUS FOR THE ENVIRONMENT

HARMFUL BY INHALATION, IN CONTACT WITH SKIN AND IF SWALLOWED.

RISK OF SERIOUS DAMAGE TO EYES.

MAY CAUSE SENSITISATION BY SKIN CONTACT.

VERY TOXIC TO AQUATIC ORGANISMS, MAY CAUSE LONG TERM ADVERSE EFFECTS IN THE AQUATIC ENVIRONMENT.

AVOID CONTACT WITH SKIN

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY WITH PLENTY OF WATER AND SEEK MEDICAL ADVICE

WEAR SUITABLE PROTECTIVE GLOVES.

WEAR EYE/FACE PROTECTION.

THIS MATERIAL AND ITS CONTAINER MUST BE DISPOSED OF IN A SAFE WAY.

USE APPROPRIATE CONTAINMENT TO AVOID ENVIRONMENTAL CONTAMINATION.

To avoid risks to man and the environment, comply with the instructions for use.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when applying by hand-held equipment.

However engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Environmental protection

Do not contaminate surface water or ditches with chemical or used container.

KEEP LIVESTOCK out of treated areas for at least two weeks following treatment and until poisonous weeds such as Ragwort have died and become unpalatable.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY and dispose of safely.

DO NOT RE-USE CONTAINER, for any purpose.